

Br

12

EUROPEAN PATENT APPLICATION

21 Application number: **85107220.7**

51 Int. Cl.⁴: **C 07 D 243/08, A 61 K 31/55**

22 Date of filing: **12.06.85**

30 Priority: **19.06.84 IT 2147184**

71 Applicant: **POLI INDUSTRIA CHIMICA S.p.A., Piazza Agrippa, 1, I-20141 Milano (IT)**

43 Date of publication of application: **02.01.86**
Bulletin 86/1


72 Inventor: **Massaroli, Gian Giacomo, Via Lauro, Rovegnate (Como) (IT)**

84 Designated Contracting States: **AT BE CH DE FR GB IT LI LU NL SE**

74 Representative: **Blanchetti, Giuseppe, Studio Consulenza Brevettuale Via Senato 24, I-20121 Milan (IT)**

54 "1-Bis-(4-fluorophenyl)methyl-4-(3-phenyl-2-propenyl)-hexahydro-1H-1,4-diazepine, preparation thereof and compositions containing it

57 The compound 1-[bis-(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)-hexahydro-1H-1,4-diazepine of formula I




is endowed with calcium antagonistic activity useful in human therapy.

- 1 -

1-BIS-(4-FLUOROPHENYL)METHYL-4-(3-PHENYL-2-PROPENYL)-
HEXAHYDRO-1H-1,4-DIAZEPINE, PREPARATION THEREOF AND COMPO-
SITIONS CONTAINING IT

The present invention refers to 1-bis-(4-fluorophenyl)methyl-4-(3-phenyl-2-propenyl)-hexahydro-1H-1,4-diazepine of formula I

5



10 and its non toxic, pharmaceutically acceptable salts.

A particularly preferred salt is the dihydrochloride of compound I, hereinafter specifically referred to in the disclosure.

A further object of the invention is provided by a
 15 process for the preparation of compound I and of its salts and by pharmaceutical compositions containing them as active principle.

The compound I is endowed with calcium antagonistic activity, which has been determined in vitro on fragments
 20 of rat's thoracic aorta depolarized by K^+ , according to its ability to inhibit the Ca^{++} induced spasms.

The compound I proved to be highly active in reducing "in vitro" the increase of the intraluminal pressure of the rat's caudal artery, induced by electrical stimuli.

25 In this test the concentration of compound I which causes a 50% increase of the intraluminal pressure turned

- 2 -

out to be comprised between 1 and 2 mcg/ml.


"In vivo", the compound according to the present invention proved to be remarkably active in decreasing the increase of the peripheric vascular resistances of the 5 cat's rear limb, induced by continuous infusion or nor-adrenaline.

The dose of I reducing of 50% the peripheric vascular resistances turned out to be comprised in this test, between 50 mg/kg and 75 mg/kg by endovenous route.

10 As a consequence, the compound I is useful in human therapy for the treatment of circulatory diseases of different kind and nature, being moreover endowed with a low toxicity.

15 The compound I may be administered in form of pharmaceutical compositions containing it or one of its salt as the active principle, alone or in combination with the conventional excipients, carriers, sweetening and flavouring agents normally used in the pharmaceutical practice.

20 The process for the preparation of the compound I, according to the present invention, is illustrated in the following scheme:



The 1-ethoxycarbonyl-hexahydro-1,4-diazepine is alkylated by a cinnamyl halide in the presence of an acid-base catalyst: the ethoxycarbonyl group is then removed by alkaline hydrolysis and the resulting compound is reacted with a 4,4'-difluorobenzhydryl halide, in the presence of bases. The compound so obtained may be then optionally salified according to known methods.

20 Suitable solvents which can be suitably used are acetone, ethyl acetate, methylisobutylketone, methylene chloride etc.

It is moreover obvious that protective groups other than the ethoxycarbonyl one may be used.

25 The following example further illustrates the invention without limiting it in any way.

EXAMPLE

a) 4-(3-Phenyl-2-propenyl)-1H-hexahydro-1,4-diazepine

39.65 Grams (0.26 moles) of cinnamyl chloride and
30 39.2 g (0.4 moles) of triethylamine were added to 34.4 g

- 4 -

(0.2 moles) of 1-ethoxycarbonyl-hexahydro-1,4-diazepine in 200 ml of methylisobutylketone. The mixture was refluxed for 3 hours, cooled and poured into water. The organic phase was separated, washed with water, the solvent was 5 evaporated and the residue distilled under reduced pressure, obtaining 51 g (88.5% yield) of a product having b.p. 183-186°/0.2 mm which was refluxed for 20 hours with 1000 ml of ethanol containing 200 g of KOH, obtaining after partial evaporation of the solvent, dilution with water, 10 extraction with toluene and distillation 25.7 g of 4-(3-phenyl-2-propenyl)-1H-hexahydro-1,4-diazepine having b.p. 142-144°/0.2 mm. Yield 59.3%.

b) 1-[Bis-(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)-hexahydro-1H-1,4-diazepine, dihydrochloride

15 A solution of 10.8 g (0.05 moles) of 4-(3-phenyl-2-propenyl)-1H-hexahydro-1,4-diazepine, 11.92 g (0.05 moles) of bis(4-fluorophenyl)-chloromethane and 14 ml (0.1 ml) of triethylamine in 50 ml of methylisobutylketone was refluxed for 6 hours. After cooling, the mixture was wa- 20 shed with water to neutrality, the organic phase was separated and dried on anhydrous sodium sulphate and treated under stirring with 0.1 ml of gaseous hydrochloric acid.

The dihydrochloride was filtered and crystallized from absolute ethanol, obtaining 18 g (73.2% yield) of 25 1-[Bis-(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)-hexahydro-1H-1,4-diazepine, dihydrochloride having melting point 175°C.

Argentometric assay: 99.7%.

Acidimetric assay (perchloric acid): 99.9%.

30 The F, N, C, H values are in agreement with the

- 5 -

reported formula I.

The compound so prepared can be administered by the oral route in form of tablets, sugar-coated tablets, capsules, drops, solution and the like.

5 The following formulations are cited by way of an example:


- a) tablets containing from 5 to 25 mg of compound I, mixed with excipients and disgregants conventionally used in the pharmaceutical practice;
- 10 b) solution for the administration in form of drops, containing from 5 to 25 mg of compound I in 1 ml of hydroalcoholic vehicle containing the conventional sweetening and flavouring agents commonly used in the pharmaceutical technique.

- 6 -

CLAIMS for the Contracting States:

BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. 1-[Bis-(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)-hexahydro-1H-1,4-diazepine of formula I



10

and salts thereof with non toxic, pharmaceutically acceptable acids.

2. 1-[Bis-(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)-hexahydro-1H-1,4-diazepine dihydrochloride.

3. A process for the preparation of the compound of formula I characterized in that 1-ethoxycarbonyl-hexahydro-1,4-diazepine is reacted with a cinnamyl halide in the presence of bases, that the ethoxycarbonyl group is removed by alkaline hydrolysis and that the obtained compounds is reacted with a 4,4'-difluoro-benzhydriyl halide, in the presence of bases.

4. A process according to claim 3 characterized by using cinnamyl chloride and bis-(4-fluorophenyl)-chloromethane in the presence of triethylamine.

5. Pharmaceutical compositions having calcium antagonistic activity containing as the active principle the compound I or one of its salts in admixture with suitable vehicles and excipients.

6. Compositions according to claim 4 characterized by

30

0166302

- 7 -

containing as active principle the dihydrochloride of compound I.


7. Oral, liquid or solid compositions containing from 5 to 25 mg of the compound I or of salts thereof.

- 8 -

CLAIMS for AT

1. A process for the preparation of the compound of formula I

5



10

characterized in that 1-ethoxycarbonyl-hexahydro-1,4-diazepine is reacted with a cinnamyl halide in the presence of bases, that the ethoxycarbonyl group is removed by alkaline hydrolysis and that the obtained compounds is
15 reacted with a 4,4'-difluoro-benzhydryl halide, in the presence of bases.

2. A process according to claim 1 characterized by using cinnamyl chloride and bis-(4-fluorophenyl)-chloromethane in the presence of triethylamine.

12

EUROPEAN PATENT APPLICATION

21 Application number: 85107220.7

51 Int. Cl.: **C 07 D 243/08, A 61 K 31/55**

22 Date of filing: 12.06.85

30 Priority: 19.06.84 IT 2147184

71 Applicant: **POLI INDUSTRIA CHIMICA S.p.A., Piazza Agrippa, 1, I-20141 Milano (IT)**

43 Date of publication of application: 02.01.86
Bulletin 86/1

72 Inventor: **Massaroli, Gian Giacomo, Via Lauro, Rovegnate (Como) (IT)**


84 Designated Contracting States: **AT BE CH DE FR GB IT LI LU NL SE**

88 Date of deferred publication of search report: 26.03.86 Bulletin 86/13

74 Representative: **Bianchetti, Giuseppe, Studio Consulenza Brevettuale Via Rossini, 8, I-20122 Milan (IT)**

54 "1-[Bis-(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)-hexahydro-1H-1,4-diazepine, preparation thereof and compositions containing it.

57 The compound 1-[bis-(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)-hexahydro-1H-1,4-diazepine of formula I



is endowed with calcium antagonistic activity useful in human therapy.

EP 0 166 302 A3



European Patent
Office

EUROPEAN SEARCH REPORT

0166302
Application number

EP 85 10 7220

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Y	FR-A-2 159 369 (JANSSEN PHARMACEUTICA N.V.) * Page 9, line 30; pages 11-12 *	1-7	C 07 D 243/08 A 61 K 31/55
Y	--- CHEMICAL ABSTRACTS, vol. 93, no. 15, 13th October 1980, page 42, abstract no. 142831r, Columbus, Ohio, US; C.K. SCOTT et al.: "The effects of flunarizine, a new calcium antagonist, on human red blood cells in vitro" & ANGIOLOGY 1980, 31(5), 320-30 -----	1-7	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 07 D 243/00 A 61 K 31/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 12-12-1985	Examiner BACON, A. J.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

